arXiv:0712.0855v1 [nucl-th] 6 Dec 2007

Few-Body Systems 0, 1–3 (2008)

Few-
Body
Systems

© by Springer-Verlag 2008
Printed in Austria

# *Ab initio* no-core shell model and microscopic reactions: recent achievements

S. Quaglioni *, P. Navrátil

Lawrence Livermore National Laboratory, L-414, P.O. Box 808, Livermore, CA 94551, USA

**Abstract.** We report on recent microscopic calculations of reaction properties based upon the nuclear structure of the *ab initio* no-core shell model (NCSM).

## 1 Introduction

We have explored two avenues for the *ab initio* calculations of nuclear-reaction properties starting from the nuclear structure of the NCSM [1]: i) the Lorentz integral transform (LIT) approach [2], and ii) the resonating-group method (RGM) [3]. In Sect. 2, we highlight a recent application of the NCSM/LIT approach to the *ab initio* calculation [4] of the $^4$He total photo-absorption cross section. In Sect. 3, we outline the NCSM/RGM approach and present its application to neutron-$^4$He scattering.

## 2 Application of chiral effective field theory two- plus three-nucleon forces to the $^4$He photodisitegration

We performed an *ab initio* calculation [4] of the $^4$He total photo absorption cross section in unretarded dipole approximation, using the high quality NN potential at the fourth order (N$^3$LO) in the chiral effective field theory ($\chi$EFT) expansion of Ref. [5], and the NNN interaction at the highest order presently available (N$^2$LO) [6]. The two low-energy constants that parameterize the short-range NNN interaction were selected as discussed in Ref. [7]. The microscopic treatment of the continuum problem is achieved by means of the LIT method, applied within the NCSM approach. Accurate convergence of the NCSM expansions is reached using three-body effective interactions for both the NN and NNN terms of the potential. The results, presented in Fig. 1, show a peak around the excitation energy of $\omega = 27.8$ MeV, with peak-height mildly sensitive to the NNN force. The experimental situation in the near-threshold region is controversial [8]: we find an overall good agreement with the photo-disintegration data from bremsstrahlung photons of Nilsson et al. and the indirect measurements of Nakayama et al.

---

*E-mail address:* quaglioni1@llnl.gov



**Figure 1.** The $^4$He photo-absorption cross section as a function of the excitation energy $\omega$. Present results with $\chi$EFT interactions compared to the most recent experiments [8].

## 3 Augmenting the NCSM by the RGM technique

For reactions involving two nuclear fragmets, the RGM many-body wave function is expanded in terms of binary-cluster channel wave functions

$$\Psi^{(A)} = \sum_\nu \hat{\mathcal{A}} \left[ \psi_{1\nu}^{(A-a)} \psi_{2\nu}^{(a)} \varphi_\nu(\boldsymbol{r}_{A-a,a}) \right] = \sum_\nu \int d\boldsymbol{r}\, \varphi_\nu(\boldsymbol{r}\,)\, \hat{\mathcal{A}}\, \Phi_{\nu\boldsymbol{r}}^{(A-a,a)} \,, \tag{1}$$

with

$$\Phi_{\nu\boldsymbol{r}}^{(A-a,a)} = \psi_{1\nu}^{(A-a)} \psi_{2\nu}^{(a)} \delta(\boldsymbol{r} - \boldsymbol{r}_{A-a,a}) \,. \tag{2}$$

Here, $\hat{\mathcal{A}}$ is the antisymmetrizer accounting for the exchanges of nucleons between the two clusters (which are already antisymmetric with respect to exchanges of internal nucleons). The relative-motion wave functions $\varphi_\nu$ depend on the relative distance between the center of masses of the two clusters in channel $\nu$. They can be determined by solving the many-body Schrödinger equation in the Hilbert space spanned by the basis functions (2):

$$H\Psi^{(A)} = E\Psi^{(A)} \longrightarrow \sum_\nu \int d\boldsymbol{r} \left[ \mathcal{H}_{\mu\nu}^{(A-a,a)}(\boldsymbol{r}\,', \boldsymbol{r}\,) - E\mathcal{N}_{\mu\nu}^{(A-a,a)}(\boldsymbol{r}\,', \boldsymbol{r}\,) \right] \varphi_\nu(\boldsymbol{r}\,) \,, \tag{3}$$

where $H$ denotes the full Hamiltonian of the system, and the two integration kernels are defined as

$$\mathcal{H}_{\mu\nu}^{(A-a,a)}(\boldsymbol{r}\,', \boldsymbol{r}\,) = \left\langle \Phi_{\mu\boldsymbol{r}'}^{(A-a,a)} \middle| \hat{\mathcal{A}}\, H\, \hat{\mathcal{A}} \middle| \Phi_{\nu\boldsymbol{r}}^{(A-a,a)} \right\rangle , \tag{4}$$

$$\mathcal{N}_{\mu\nu}^{(A-a,a)}(\boldsymbol{r}\,', \boldsymbol{r}\,) = \left\langle \Phi_{\mu\boldsymbol{r}'}^{(A-a,a)} \middle| \hat{\mathcal{A}}^2 \middle| \Phi_{\nu\boldsymbol{r}}^{(A-a,a)} \right\rangle . \tag{5}$$

In the NCSM/RGM approach we use *ab initio* NCSM wave functions for each of the two clusters, and NCSM effective interactions derived from realistic forces. The most challenging task is to evaluate the Hamiltonian and norm kernels: so far we have developed formalism for cluster states with a single-nucleon projectile and we have applied it to the scattering of low-energy neutrons on $^4$He. In Fig. 2,



**Figure 2.** The calculated $^2S_{1/2}$ phase shift for the $n$+$^4$He system as a function of the relative kinetic energy $E$, compared to experiment. The $V_{lowk}$ [9] potential at $\hbar\Omega = 18$ MeV in a $16\hbar\Omega$ model space, and the $\chi$EFT NN interaction [5] at $\hbar\Omega = 19$ MeV in a $14\hbar\Omega$ model space were used.

we present the converged $^2S_{1/2}$ phase shift for the $n$+$^4$He system obtained using the bare $V_{lowk}$ [9] potential, and two-body effective interactions derived from the $\chi$EFT NN potential of Ref. [5]. We find an overall good agreement with experiment.

*Acknowledgement.* This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344. Support from U.S. DOE/SC/NP (Work Proposal Number SCW0498) and the Department of Energy under Grant DE-FC02-07ER41457 is acknowledged.

## References

1. Navrátil, P., et al.: Phys. Rev. Lett. **84**, 5728 (2000); Phys. Rev. **C62**, 054311 (2000); Phys. Rev. Lett. **88**, 152502 (2002); Phys. Rev. **C68**, 034305 (2003)
2. Efros, V. D., et al.: Phys. Lett. **B338**, 130 (1994)
3. Tang, Y. C., et al: Phys. Rep. **47**, 167 (1978); Langanke, K., Friedrich, H.: Advances in Nuclear physics. NewYork: Plenum (1987)
4. Quaglioni, S., Navrátil, P.: Phys. Lett. **B652**, 370 (2007)
5. Entem, D. R., Machleidt, R.: Phys. Rev. **C68**, 041001(R) (2003)
6. Van Kolck, U.: Phys. Rev. **C49**, 2932 (1994); Epelbaum, E., et al.: Phys. Rev. **C66**, 064001 (2002)
7. Navrátil, P., et al.: Phys. Rev. Lett. **99**, 042501 (2007)
8. Shima, T., et al.: Phys. Rev. **C72**, 044004 (2005); Nillson, B., et al.: Phys. Rev. **C75**, 014007 (2007); Nakayama, S., et al.: Phys. Rev. C76, 021305(R) (2007)
9. Bogner, S. K., Kuo, T. T. S., Schwenk, A.: Phys. Rep. **386**, 1 (2003)